
(12) UK Patent Application (19) GB (11) 2 080 524 A

(21) Application No 8119357

(22) Date of filing 23 Jun 1981

(30) Priority data

(31) 163456

(32) 27 Jun 1980

(31) 254233

(32) 14 Apr 1981

(33) United States of America
(US)

(43) Application published
3 Feb 1982

(51) INT CL³
G01N 33/48

(52) Domestic classification
G1B BV

(56) Documents cited
GB 2033081A
GB 2019563A
GB 1530810

(58) Field of search
G1B

(71) Applicant
Mallinckrodt, Inc.,
675 McDonnell Blvd., P.O.
Box 5840, St. Louis,
Missouri 63134, United
States of America

(72) Inventor
Roxane N. Dikeman

(74) Agents
Batchellor, Kirk & Eyles,
2 Pear Tree Court,
Farringdon Road, London,
EC1R 0DS

(54) Determining Endotoxin with a
Limulus Amebocyte Lysate Reagent

(57) A reagent, for determining
endotoxin, comprises a buffered
aqueous dispersion of Limulus
amebocyte lysate and a surfactant as
a lysate sensitivity enhancer.

GB 2 080 524 A

SPECIFICATION

Improvement in the Process for Preparing Limulus Lysate

This invention relates to a process for improving the sensitivity of Limulus amebocyte lysate (hereinafter sometimes referred to as LAL or lysate) to endotoxin, to an improved LAL reagent and to the use of such LAL reagent.

As is well known, the LAL test for detecting endotoxins is perhaps the most practical and sensitive test for determining endotoxins. Commercial assay tests employ amebocyte lysate from Limulus hemolymph obtained from the horseshoe crabs. This lysate is combined with appropriate divalent cations, appropriate buffers and other ingredients to form a LAL reagent. This reagent then reacts with endotoxin during the assay to form a gel. Manufacturers of LAL reagents often experience difficulty in producing lysate of the desired sensitivity for detecting endotoxin. Sensitivity from one preparation to the next is also variable. These problems are attributed at least in part to the presence of an endogenous, undefined endotoxin inhibitor substance in the lysate, (hereinafter sometimes referred to as the inhibitor).

Little is known of the nature of the inhibitor or its in-vivo role in the horseshoe crab. Electrophoretic studies indicate that the inhibitor is a high molecular weight lipoprotein. It may function in the amebocyte to control the coagulation defense mechanism. That the inhibitor may be a membrane component freed during cell lysis is also plausible. The uncertainty of the role and origin of the inhibitor is compounded by the fact that the mechanism of inhibition is unclear. The inhibitor presumably blocks the enzymatic reaction in some fashion either by association with the enzyme itself, or with the endotoxin or both. Like some other serine proteases, the proclotting enzyme is thought to be complexed with calcium and glycerophospholipid. Endotoxin itself is lipoidal, hence, an inhibitor of lipoprotein character would be highly compatible with either component.

That the inhibitor is a lipoprotein is supported by its sensitivity to chloroform. As described in U.S. Patent No. 4,107,077, the sensitivity of LAL is improved substantially when lysate is treated with an organic solvent such as chloroform to precipitate inhibitor from the lysate. The aqueous phase is then recovered and processed to prepare the LAL reagent.


To date, the above-mentioned solvent extraction procedure is the most rapid means of improving LAL sensitivity. Unfortunately, the method has several drawbacks. Because of the absolute requirement that endotoxin-free conditions be maintained throughout the lysate production, a cumbersome extraction procedure and subsequent centrifugation increases the likelihood of product failure. As noted in the patent, the solvent treatment reduces lysate stability such that the production must be completed rapidly in the cold. Also the precipitate removed from the lysate by the solvent treatment contains considerable coagulogen, the required clotting protein. Maintenance of adequate protein content is a requirement for firm gelation during endotoxin assay. Obviously, under the latter circumstances, control of reagent sensitivity is difficult. Chloroform, the solvent used most successfully, is well known for its undesirable effects in man. The health and safety of production personnel is therefore a reasonable concern. It is apparent that a process which avoids these pitfalls and yet improves sensitivity to the desired degree would be an improvement in the art.

It is therefore an object of the present invention to provide a method for the simple and rapid enhancement of the sensitivity of LAL.

In accordance with this invention, there is provided a process for treating under lysate treating conditions LAL having decreased sensitivity to endotoxin due to the presence of an endogenous inhibitor with an enhancing amount of a lysate sensitivity enhancing agent to neutralize or partially neutralize the lysate inhibitor thereby increasing the LAL sensitivity to endotoxin.

There are certain minimal criteria which can delineate LAL sensitivity enhancing characteristics, i.e., the LAL sensitivity enhancing agents useful in the process of this invention should possess (a) the ability to increase the lysate sensitivity to a suitable sensitivity, e.g., by a twofold or greater increase, (b) the ability to withstand depyrogenation, i.e. removal or destruction of endotoxin, by ultrafiltration or acid treatment at pH less than 5 or base treatment greater than pH 8, (c) the ability to be sterilized e.g., by autoclaving e.g. at or above 121°C at 15 psi for 15 minutes, (d) the ability to form aqueous solutions of about 2% (w/v) at 25°C, (e) the ability to function in the pH range of about 6.0 to about 9, (f) the ability to be compatible with buffers and other ingredients utilized in the LAL reagent and (g) the ability to be compatible with respect to LAL and its reaction with endotoxins.

Such enhancing agents include amphoteric surfactants which have both an anionic and cationic group in their structure. Illustrative are the sulfobetaines represented by the following formula (hereinafter Formula A:



wherein:

R_1 is an alkylene radical having from 1 to about 4 carbon atoms,

Y is any non-deleterious, chemically suitable substituent including (1) hydrogen, (2) substituted or unsubstituted lower alkyl, e.g. containing 1 to 4 carbon atoms such as methyl, ethyl, propyl, or hydroxy ethyl, hydroxy methyl, hydroxy propyl, etc.;

R_2 and R_3 are each selected from substituted or unsubstituted lower alkyl containing 1 to 4 carbon atoms, e.g., such as methyl, ethyl, propyl, hydroxy ethyl, hydroxy methyl, hydroxy propyl, etc. $n=0$ or 1,

when $n=0$, R_4 is substituted or unsubstituted alkyl, e.g. containing about 8 to about 18 carbon atoms,

when $n=1$, R_4 is an alkylene radical having from about 1 to about 6 carbon atoms,

R_5 is a substituted or unsubstituted alkyl, e.g. containing about 8 to about 18 carbon atoms;

It is to be understood that the term "alkylene" as it is used herein, encompasses both polymethylene radicals and other divalent saturated aliphatic radicals and thus there may be branching in the linkage provided by the alkylene radical. The item "lower" means a radical containing 1 to 4 carbon atoms.

The sulfobetaines which are employed in the compositions of the present invention are known in the art and have been described as zwitterionic surfactants. The preparation of such compounds is described, for example, by G. W. Fernley in the Journal of American Oil Chemists Society, January 1978 (Vol. 55), pages 98—103, and by R. Ernst in the U.S. Patent 3,280,179 issued October 18, 1966, which patent is incorporated herein by reference.

In preferred sulfobetaine surfactants, R_2 and R_3 in the above structure are methyl. It is also preferred that R_1 be propylene.

One type of sulfobetaine surfactant which can be employed has the above structure wherein n equals 0 and R_4 is an alkyl radical having from about 8 to 18 carbon atoms, preferably a straight chain alkyl radical. For these sulfobetaine surfactants, a convenient source of the R_4 component is tallow fatty alcohol which consists of a mixture of various chain lengths, with a typical composition being approximately 66 percent C_{18} , 30 percent C_{16} and 4 percent C_{14} and others. Another convenient source is the middle cut of distilled coconut fatty alcohol, which also consists of a mixture of various chain lengths, with a typical composition being approximately 66 percent C_{12} , 23 percent C_{14} , 9 percent C_{16} and 2 percent C_{10} .


Specific sulfobetaine surfactants of the above structure wherein n equals 0 are set forth in U.S. Patent 3,539,521 issued on November 10, 1970 to A. O. Snoddy et al, which patent is herein incorporated by reference. A surfactant of this type particularly preferred is N-tetradecyl-N,N-dimethyl-3-ammonio-1-propanesulfonate commercially available from Calbiochem-Behring Corporation under the trademark Zwittergent 3-14.

Another type of sulfobetaine surfactant which can be employed has the above structure wherein n equals 1 and R_4 is an alkylene radical having from about 1 to about 6 carbon atoms. In these sulfobetaines wherein n equals 1, R_5 is an alkyl radical having from about 8 to about 18 carbon atoms. It is preferred that R_5 be straight chain. As previously discussed, convenient sources of alkyl radicals having from about 10 to about 18 carbon atoms are tallow fatty alcohol and coconut fatty alcohol.


Specific sulfobetaine surfactants of the above structure wherein n equals 1 are set forth in the previously mentioned U.S. Patent 3,280,179.

Particularly preferred sulfobetaine surfactants for use in compositions of the present invention are 3-(N,N-dimethyl-N-acylamidopropylammonio)-2-hydroxypropane-1-sulfonates wherein the acyl group is derived from tallow fatty alcohol or coconut fatty alcohol, with coconut fatty alcohol preferred. It would be recognized by those skilled in the art that in the normal preparation of these derivatives of tallow or coconut fatty alcohols, a mixture of sulfobetaines with varying carbon chain lengths for the acyl groups would result. As previously discussed, these fatty alcohols contain for the most part carbon chain lengths which will provide acyl groups with the desired number of carbon atoms, that is from about 8 to about 18 carbon atoms. Thus, these mixtures obtained from tallow or coconut fatty alcohols are useful in providing the sulfobetaine surfactant in the compositions of the present invention. A material of this type particularly preferred for use in the composition of the present invention is N-cocoamido-propyl-N,N-dimethyl-N-2-hydroxypropyl sulfobetaine, an example of which is LLONZAINE CS, commercially available from Lonza, Inc., Fair Lawn, New Jersey, another example of which is VARION CAS commercially available from Sherex Chemical Company, Inc.


Other amphoteric surfactants include, the N-long chain alkyl aminocarboxylic acids illustrated by the formula (hereafter Formula B):



the N-long chain alkyl iminodicarboxylic acids illustrated by the formula (hereinafter Formula C):



and the N-long chain alkyl or amido betaines illustrated by the formula (hereinafter Formula D):




where R_1 , R_2 , R_3 , R_4 , Y and n have the same meaning as they have in Formula A, M is hydrogen or a salt-forming metal, such as sodium or potassium, or ammonium, and Y' has the same meaning as Y in Formula A. Y and Y' may be the same or different. Examples of specific amphoteric detergents are N-alkyl-beta-aminopropionic acid, N-alkyl-beta-iminodipropionic acid, and N-alkyl-N,N-dimethyl glycine; the alkyl group may be, for example, that derived from coco fatty alcohol, lauryl alcohol, myristyl alcohol (or a lauryl-myristyl mixture), hydrogenated tallow alcohol, cetyl, stearyl, or blends of such alcohols. The substituted aminopropionic and iminodipropionic acids are often supplied in the sodium or other salt forms, which may likewise be used in the practice of this invention. Specific examples include cocobetaine sold by Witco Chemical Corporation under the name EMCOL CC 37-18; cocoamidopropyl betaine sold by Lonza Inc. and Sherex Chemical Company under the names LONZAINE CO and VARION CADG, respectively; sodium N-coco-beta-aminopropionate sold by Henkel Corporation under the name DERIPHAT 151; disodium N-lauryl-beta-iminodipropionate sold by Henkel Corporation under the name DERIPHAT 160, and disodium N-tallow-beta-iminodipropionate sold by Henkel Corporation under the name of DERIPHAT 154.

Examples of other amphoteric detergents are the fatty imidazolines such as those made by reacting a long chain fatty acid (e.g. of 10 to 20 carbon atoms) with diethylene triamine and monohalocarboxylic acids having 2 to 6 carbon atoms, e.g. 1-coco-5-hydroxyethyl-5-carboxymethylimidazoline.

Specific examples include cocoimidazoline commercially available under the name AMPHOTERGE K-2 from Lonza, Inc., capric dicarboxy imidazoline commercially available under the name AMPHOTERGE KJ2 from Lonza, Inc. and coco dicarboxy imidazoline blended with sulfated surfactants commercially available under the name AMPHOTERGE 2 WAS MOD from Lonza, Inc.

Other examples of enhancing agents include anionic synthetic surfactants, generally described as those compounds which contain hydrophilic and lipophilic groups in their molecular structure and ionize in an aqueous medium to give anions containing both the lipophilic group and hydrophilic group. The alkyl aryl sulfonates, the alkane sulfates and sulfated oxyethylated alkyl phenols are illustrative of the anionic type of surface active compounds.


The alkyl aryl sulfonates are a class of synthetic anionic surface active agents represented by the general formula (hereinafter Formula E):



R_6 is a straight or branched chain hydrocarbon radical having from about 1 to about 24 carbon atoms, at least one R_6 having at least 8 carbon atoms; n_1 is from 1 to 3; n_2 is from 1 to 2; Ar is a phenyl or a naphthyl radical and Y and M have the same meaning as in Formula B. R_6 can be, for example, methyl, ethyl, hexyl, octyl, tetradecyl, iso-octyl, nonyl, decyl, dodecyl, octadecyl and the like.

Compounds illustrative of the alkyl aryl sulfonates include sodium dodecylbenzene sulfonate, sodium decylbenzene sulfonate, ammonium methyl dodecylbenzene sulfonate, ammonium dodecylbenzene sulfonate, sodium octadecylbenzene sulfonate, sodium nonylbenzene sulfonate, sodium dodecyl-naphthalene sulfonate, sodium tetradecylbenzene sulfonate, potassium eicosyl-naphthalene sulfonate, ethylamine undecyl-naphthalene sulfonate and sodium docosyl-naphthalene sulfonate.


The alkyl sulfates are a class of synthetic anionic surface active agents represented by the general formula (hereinafter Formula F):



where R_5 and M have the same meaning as in Formula B.

Compounds illustrative of alkyl sulfate class of anionic surfactants include sodium octadecyl sulfate, sodium hexadecyl sulfate, sodium dodecyl sulfate, sodium nonyl sulfate, ammonium decyl sulfate, potassium tetradecyl sulfate, diethanolamino octyl sulfate, triethanolamine octadecyl sulfate and ammonium nonyl sulfate.

The sulfated oxyethylated alkylphenols are a class of synthetic anionic surface active agents represented by the general formula (hereinafter Formula G):



where A is either oxygen, sulfur, a carbonamide group, thiocarbonamide group, a carboxylic group or thiocarboxylic ester group, Z is an integer from 3 to 8 and R_5 and M have the same meaning as in Formula B.

5 Compounds illustrative of the sulfated oxyethylated alkyl phenol class of anionic surfactants include ammonium nonylphenoxyl tetraethylenoxy sulfate, sodium dodecylphenoxyl triethylenoxy sulfate, ethanolamine decylphenoxyl tetraethylenoxy sulfate and potassium octylphenoxyl triethylenoxy sulfate.

10 Other examples of LAL enhancing agents include nonionic surface active compounds which can be broadly described as compounds which do not ionize but acquire hydrophilic characteristics from an oxygenated side chain such as polyoxyethylene and the lipophilic part of the molecule may come from fatty acids, phenol, alcohols, amides or amines. The compounds are usually made by reacting an alkylene oxide such as ethylene oxide, butylene oxide, propylene oxide and the like, with fatty acids, straight or branched chain alcohols containing one or more hydroxyl groups, phenol, thiophenols, 15 amides and amines to form polyoxyalkylene glycoethers and esters, polyoxyalkylene alkylphenols, polyoxyalkylene thiophenols, polyoxyalkylene amides and the like. It is generally preferred to react from about 3 to about 30, more preferably 10 to 30, moles of alkylene oxide per mole of the fatty acids, alcohols, phenols, thiophenols, amides or amines.

20 Illustrative of these nonionic surfactants are the products obtained from the reaction of alkylene oxide with an aliphatic alcohol having from 8 to 18 carbon atoms, such as octyl, nonyl, decyl, octadecyl, dodecyl, tetradecyl and the like; with monoesters of hexahydric alcohols containing, for example 6 carbon atoms, the ester group containing 10 to 20 carbon atoms such as sorbitan monolaurate, sorbitan monooleate and sorbitan monopalmitate; with an alkyl phenol in which the alkyl group contains between 4 and 20 carbon atoms, such as butyl, dibutyl, amyl, octyl, dodecyl, tetradecyl 25 and the like; and with an alkyl amine in which the alkyl group contains between 1 to 8 carbon atoms.


Compounds illustrative of synthetic nonionic surfactants include the products obtained from condensing ethylene oxide or propylene oxide with the following: propylene glycol, ethylene diamine, diethylene glycol, dodecyl phenol, nonyl phenol, tetradecyl alcohol, N-octadecyl diethanolamine, N-dodecyl monoethanolamide, polyoxyethylene (20) sorbitan monooleate sold under the name TWEEN 30 80 and polyoxyethylene (20) sorbitan monolaurate sold under the name TWEEN 20.

Other nonionic surfactants include long chain tertiary amine oxides corresponding to the following general formula (hereinafter Formula H):



35 wherein R_5 has the same meaning as in Formula A, and R_7 and R_8 are each methyl or ethyl radicals. The arrow in the formula is a conventional representation of a semi-polar bond. Examples of amine oxides suitable for use in this invention include dimethyldodecylamine oxide, dimethyloctylamine oxide, dimethyldodecylamine oxide, dimethyltridecylamine oxide, dimethylhexadecylamine oxide.

40 Cationic surface active agents may also be employed as LAL enhancing agents. Such agents are those surface active compounds which contain an organic hydrophobic group and a cationic solubilizing group. Typical cationic solubilizing groups are amine and quaternary groups. Such cationic surface active agents are represented by the following general formula (hereinafter Formula I)



wherein R_5 , Y and Y' have the same meaning as in Formula C. An example is QUATERNARY O available from Ciba-Geigy Corporation.

45 Other examples of suitable synthetic cationic surfactants include the diamines such as those of the formula (hereinafter Formula J):




wherein R_9 is an alkyl group of about 12 to 22 carbon atoms, such as N-2-aminoethyl stearyl amine and N-2-aminoethyl myristyl amine; amide-linked amines such as those of the formula (hereinafter

50 Formula K):



such as N-2-amino ethylstearyl amide and N-amino ethyl myristyl amide; quaternary ammonium compounds wherein typically one of the groups linked to the nitrogen atom are alkyl groups which contain 1 to 3 carbon atoms, including such 1 to 3 carbon alkyl groups bearing inert substituents, such as phenyl groups and there is present an anion such as halogen, acetate, methylsulfate, etc. Typical quaternary ammonium compounds are ethyl-dimethyl-stearyl ammonium chloride, benzyl-dimethyl-stearyl ammonium chloride, benzyl-dimethyl-stearyl ammonium chloride, trimethyl stearyl ammonium chloride, trimethylcetyl ammonium bromide, dimethylethyl dilaurylammonium chloride, dimethyl-propyl-myristyl ammonium chloride, and the corresponding methosulfates and acetates.

Another suitable cationic surfactant is represented by the formula (hereinafter Formula L):



wherein R_5 has the same meaning as in Formula A and each a is an integer from 1 to 15. An example is the polyethylene glycol amine of hydrogenated tallow wherein R_5 represents the tallow radical and a has an average value of 5. It is available from Ciba-Geigy Corporation under the trade name BINA COBA 3001.

As mentioned, the lysate enhancing agent is used in enhancing amounts, i.e. sufficient to neutralize or partially neutralize the endogenous endotoxin inhibitor in the lysate. Generally, this is an amount from about 0.001 to 1.0% (w/v) preferably from about 0.01% to about 0.05% (w/v) based on the total volume of the lysate. Frequently, amounts in excess of an enhancing amount interfere with the ability of the LAL to react with endotoxin during assay.

LAL may be prepared by those procedures known in the art, e.g. the procedure, described in British Patent 1,522,127 which is incorporated herein by reference.

For example, the hemolymph from healthy specimens of *Limulus polyphemus* is collected in a saline anticoagulant solution generally as described by Levin and Bang—"Clottable Protein in Limulus: Its Localization and Kinetics of its Coagulation by Endotoxin", *Thromb. Diath. Haemorrh.* 19: 186—197 (1968).—The amebocytes are collected and washed with the saline anticoagulant solution with the amebocyte separated from the anticoagulant by centrifugation.

The separated amebocytes are suspended in water and the osmotic disruption of the cells is complemented by mechanical agitation. The cellular debris is separated from the lysate by centrifugation and the lysate fractions are pooled and stored at 0—4°C.

To form the LAL reagent, the aforementioned LAL fractions are generally buffered to a suitable pH range, e.g. 5.5 to 8.5, preferably 6.5 to 7.5 by means of a suitable buffer, e.g. tris(hydroxymethyl)aminomethane, tris(hydroxymethyl)aminomethane maleate, 1,4-piperazine-diethanesulfonic acid, morpholinopropanesulfonic acid, N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid, triethanolamine, imidazole and tris(hydroxymethyl)imidazole. Then, the LAL reagent can be subdivided into serum vials, e.g. containing 1.2 or 5.2 ml. of solution and lyophilized. Normally, after lyophilization the vials are sealed and refrigerated (1—5°C.).

Normally, in accordance with this invention the LAL is treated by adding the lysate sensitivity enhancing agent to the LAL after the LAL has been separated from the amebocyte cellular debris. Usually, it is added prior to or at the time of preparing the LAL reagent, e.g., simultaneously with the buffer and other ingredients.

Sensitivity of the LAL reagent toward endotoxin is further increased by including low concentrations of divalent and monovalent cations. Calcium and manganese ions are the preferred divalent ions, although other alkaline earth ions such as magnesium and strontium ions or other divalent ions may be used. Magnesium and strontium ions are also preferred divalent ions. Sodium ions are the preferred monovalent ions, but other monovalent ions, especially alkali metal ions such as lithium ions may be used. The chlorides (CaCl_2 , NaCl , etc.) are convenient sources of these added ions, although other salts may be used. Preferably these electrolytes are added in endotoxin sensitivity increasing amounts, e.g. for the divalent cation (e.g., Ca^{+2}), the concentration will be in the range of 0.0001—0.4 molar and for the monovalent cation (e.g., Na^{+}), the concentration will be in the range of 0.01—0.4 molar.

The LAL reagent may also contain conventional adjuvants such as stabilizers, including lactose. These adjuvants when employed are provided in minor amounts sufficient to impart the intended qualities, but not adverse to, the desired properties of the LAL reagents.

All of the above operations are carried out under lysate treating conditions, which include insuring that the final product is sterile and free of endotoxin. Methods of insuring freedom from

of their melting points, etc., must ordinarily be dissolved, rendered acidic (pH<5) or alkaline (pH>9), and the solution autoclaved at 121°C for 30—60 minutes or more to destroy any endotoxins present.

As mentioned, another aspect of the invention is directed to a LAL reagent containing as the essential ingredient an aqueous dispersion of LAL, a LAL sensitivity enhancing agent as described previously in a lysate enhancing amount and a suitable buffer described previously in a buffering amount. Optionally, monovalent and divalent cations described above can be included in lysate sensitivity increasing amounts to further increase the sensitivity of lysate to endotoxin.

Normally, the lysate in the reagent of this invention is present in an endotoxin determining amount, e.g. an amount sufficient to determine endotoxins in a subsequent LAL assay for endotoxins, and generally this is an amount that will detect about 0.007 to about 0.5 ng/ml, preferably from about 0.007 to about 0.050 ng/ml of FDA Reference Endotoxins EC-2. The aforementioned LAL reagent can be lyophilized which is preferred.

In accordance with this invention the LAL reagent can be utilized to determine endotoxin under endotoxin determining conditions according to the usual procedure, e.g. as described in British Patent 1,522,127, and hereafter in the Examples.

The following examples illustrate the invention; all parts are by weight/volume unless otherwise stated.

Preparation of Limulus Lysate

Limulus Lysate was prepared by modification of a procedure described originally by Levin and Bang (Thromb. Diath. Haemorrh. 19, 186 (1969). Horseshoe crabs, *Limulus polyphemus*, were taken from the Atlantic Ocean in the vicinity of Beaufort, N. C. Hemolymph (approximately 500 ml) removed by cardiac puncture with a 16-gauge needle was collected in an endotoxin-free one liter glass centrifuge bottle which contained 500 ml of 0.125% N-ethylmaleimide in endotoxin-free 3% saline warmed to 42°C. The centrifuge bottle containing hemolymph-anticoagulant solution was warmed to 42° for 8 minutes and then centrifuged at 150xg for 10 minutes. The plasma supernatant was decanted and the amebocyte pellet was resuspended in the anticoagulant solution. The cells were again pelleted by centrifugation as before. The packed cells were resuspended in 0.9% pyrogen-free saline and transferred to a depyrogenated 50 ml plastic centrifuge tube. The washed cells were centrifuged again at 150xg. After decanting the saline, the packed amebocytes were ruptured by addition of pyrogen-free water for injection in a ratio of 7 ml water to 3 ml packed cells. After mixing on a vortex for 10—15 seconds. The lysed cells were stored for 24 hours at 1—5°C. Cell debris was sedimented by centrifugation at 1500xg for approximately 15 minutes. The lysate was decanted and stored at 0—4°C. The cell debris was discarded.

Preparation of Standard Endotoxin Solutions

Standard solutions of Food and Drug Administration (FDA) reference standard endotoxin Lot EC-2 were prepared in pyrogen-free water for injection. Reconstitution of 1 µg endotoxin supplied in a vial with 10 ml water resulted in an initial concentration of 0.1 µg/ml. The vial was shaken on a reciprocal shaker for 1 hour. Serial dilutions were prepared to provide the following endotoxin concentrations: 10 ng/ml, 1 ng/ml, 500 pg/ml, 250 pg/ml, 125 pg/ml, 62.5 pg/ml, 31.25 pg/ml, 15.6 pg/ml, and 7.8 pg/ml. Once prepared, endotoxin solutions were stored up to 48 hours and then discarded. Other endotoxin standards used were solutions of FDA reference Endotoxin Lot No. 1 from *Klebsiella pneumoniae*, *Escherichia coli* endotoxin Lot 071857 (Difco), and a reformulation of reference Lot EC-2 prepared in our laboratory. Endotoxin standard solutions prepared from *E. coli* Lot 071857 endotoxin were of the following concentrations: 6.25, 12.5, 50, 75, 100, 150, and 200 pg/ml.

Lysate Assay Procedure

Lysate dilutions of 25 to 70% were prepared in a 0.1M buffer pH 7.0 which was usually tris, i.e., tris (hydroxymethyl)aminomethane. Other buffers used included imidazole, tris imidazole, triethanolamine, tris maleate, N-2-hydroxyethylpiperazine-N-2-ethanesulfonic acid (HEPES), 1,4-piperazinediethane sulfonic acid (PIPES), and morpholinopropane sulfonic acid (MOPS). To determine the sensitivity of the lysate, 0.1 ml of each of the endotoxin dilutions was combined with 0.1 ml of lysate in depyrogenated 10x75 mm screw capped glass tubes and incubated for 1 hour at 37°C. Results were determined by gently inverting each tube to 180°. A clot which remained intact after the inversion indicated a positive endotoxin test.

Preparation and Depyrogenation of Surfactant Solutions

Stock LAL sensitivity enhancing agent solutions were prepared in concentrations up to 10% active ingredients in aqueous solution (w/v). Most frequently, the concentration prepared was 1% (w/v). Although the procedure varied in amounts from one agent to the next, enough material was dissolved in 50 ml aqueous solution to yield the desired concentration if diluted to 100 ml. The solution also contained 7.5 ml of 0.05M tris(hydroxymethyl)aminomethane (i.e. TRIZMA BASE, Sigma Chemical Company) and 1 ml of 2N NaOH to give a final pH≥11. The agent in alkaline solution was stored for 12 hours or more at 0—4° to insure complete denaturation. After adjusting the solution

to approximately pH 8, it was autoclaved at $\geq 121^{\circ}\text{C}$ at 15 psi for 15 minutes or more. The pH was adjusted finally to $\text{pH } 7.0 \pm 0.5$. The agent concentration was calculated and the dilution adjusted to yield the final desired concentration. Alkali-labile agents were depyrogenated by acid treatment in which HCl and tris buffer were substituted for NaOH and TRIZMA BASE, respectively.

5 Addition of Agents to Lysate 5

Agent solutions prepared as described above were added to the lysate during dilution with buffer. The amount added varied with the agent used, but the concentration range for all those tested was 0.001 to 1.0% (w/v) final concentration in the lysate solution. The order of addition of components did not alter the resulting lysate sensitivity.

10 Example I 10

The LAL sensitivity enhancing agent, ZWITTERGENT™ 3—14, N,N-dimethyl-3-ammonio-1-propanesulfonate, a sulfobetaine sold by Calbiochem-Behring Corporation, was depyrogenated as described above and diluted to a final 1% concentration (w/v). Lysate lot 9C2C was prepared as a 50% dilution with 0.1M tris-maleate buffer pH 7.0 addition of the enhancing agent was carried out to yield 15 final concentration in lysate of 0.005, 0.01, 0.02, 0.05, 0.075, and 0.10% (w/v). A control sample 15 contained pyrogen-free water instead of surfactant. The lysate dilutions were stored overnight at $0-4^{\circ}\text{C}$ and tested the following day with a specially formulated EC endotoxin series of dilutions (designated EC). The results (Table I) indicate both the effective concentration range for enhancing agent and the total increase in lysate sensitivity to endotoxin as compare to the control lysate.

Table I 20

Sample	Enhancing Agent (%)	Lysate Sensitivity*
Control	0	500
LAL plus	0.005	1,000.0
Enhancing Agent	0.01	500.0
	0.02	31.2
	0.05	10,000.0
	0.075	10,000.0
	0.10	10,000.0

*Expressed as the lowest endotoxin concentration (pg/ml) which yields a positive clot test. 30

Example II

Standard endotoxin solutions were prepared for *E. coli* endotoxin 071857, FDA Reference Endotoxin EC-2, and *K. pneumoniae* FDA reference Lot No. 1. Lysate was diluted with 0.1M tris buffer pH 7.0 and the enhancing agent, ZWITTERGENT™ 3—14, solution to yield a 30% lysate dilution 35 containing 0.02% ZWITTERGENT 3—14 (w/v). The agent was replaced by water in the control sample. 35 The lysate assay was carried out with each endotoxin dilution series.

Table II

Endotoxin	Sample	Lysate Sensitivity*
<i>E. coli</i> 071857	Control	75.0
	LAL enhancing agent	25.0
	Control	500.0
	LAL enhancing agent	62.5
FDA Reference EC-2	Control	1,000.0
	LAL enhancing agent	125.0

*Expressed as the lowest endotoxin concentration (pg/ml) which yields a positive clot test.

Example III

Twenty-two commercially available LAL enhancing agents were surveyed to determine their effects on lysate sensitivity to endotoxin. Solutions of each were prepared and added to lysate Lot 9FI 50 in final concentrations ranging from 0.001 to 0.20% (w/v). Lysate samples were then tested with reformulated FDA EC endotoxin (EC). In Table III, the agents are listed in order of decreasing effectiveness. The most effective concentration tested in lysate and its corresponding lysate sensitivity 50 are shown. The sensitivity of control lysate (50% dilution) containing no agent is also indicated.

Table III

	Trade Name	Source	Surfactant	Effective Concentration (%)	^a Lysate Sensitivity
	Zwittergent 3—14	Calbiochem-Behring	Amphoteric	0.02	15.6
5	Varion CADG	Sherex Chemical Company, Inc.	Amphoteric	0.02	15.6
	Lonzaine CO	Lonza Inc.	Amphoteric	0.02	15.6
	Lonzaine CS	Lonza Inc.	Amphoteric	0.02	15.6
	Varion CAS	Sherex Chemical Company, Inc.	Amphoteric	0.02	31.2
	Emcol CC38-17	Witco Chemical Corporation	Amphoteric	0.02	31.2
10	Deriphat 151	Henkel Inc. USA	^b Amphoteric	0.01	31.2
	Bina Coba 3001	Ciba-Geigy Corporation	Cationic	0.01	31.2
	Barlox 12	Lonza Inc.	Nonionic	0.01	31.2
	Quaternary O	Ciba-Geigy Corporation	Cationic	0.002	31.2
	Polystep B-5	Stepan Chemical Company	Anionic	0.001	31.2
15	Tween 80	ICI Americas Inc.	Nonionic	0.20	62.5
	Amphoterge K-2	Lonza Inc.	Amphoteric	0.02	62.5
	Lodyne S-100	Ciba-Geigy Corporation	Amphoteric	0.02	62.5
	Deriphat 154	Henkel Inc. USA	^b Amphoteric	0.002	62.5
	Polystep A-15	Stepan Chemical Company	Anionic	0.002	62.5
20	Amphoterge KJ-2	Lonza Inc.	Amphoteric	0.001	62.5
	Deriphat 160	Henkel Inc. USA	^b Amphoteric	0.02	62.5
	Polystep A-17	Stepan Chemical Company	Anionic	0.001	62.5
	Amphoterge WAS	Lonza Inc.	Amphoteric	0.001	125.0
	Dianol RSX	Quaker Chemical Corporation	Anionic	0.02	250.0
25	Tween 20	ICI Americas Inc.	Nonionic	0.002	250.0
	Control Lysate	—	—	—	125.0

^aExpressed as the lowest endotoxin concentration (pg/ml) which yields a positive clot test.

^bAt pH 7, the anionic form predominates.

Example IV

- 30 To determine that the increased sensitivity observed in lysate treated with the enhancing agent 30
was maintained during lyophilization, 30% dilution of lysate in 0.05M tris buffer with and without
0.02% ZWITTERGENT™ 3—14 (final concentration in lysate) were prepared. Lysate solution (1.2 ml)
was dispensed into each 10-ml serum vial. Samples were frozen at -35°C and lyophilized under 50 μ
vacuum with a drying time of approximately 32 hours. The vials were sealed with split rubber stoppers
35 and metal caps. The freeze-dried lysate was reconstituted with 1.2 ml pyrogen-free water for injection 35
and then tested with FDA Reference Endotoxin lot EC-2. Control lysate without the enhancing agent
had a sensitivity of 62.5 pg/ml. In the presence of the enhancing agent, the sensitivity of the lysate was
improved by two fold to 31.2 pg/ml.

Example V


- 40 Limulus Lysate Day Pool Lot ODH approximately one week old was prepared as a 40% dilution in 40
0.05M tris(hydroxymethyl) aminomethane maleate buffer, pH 7.0, containing in final concentrations
0.06M CaCl₂ and 0.01M MnCl₂ and 0.03% ZWITTERGENT™ 3—14. This solution was dispensed as
1.2 ml aliquots into 8 ml vials and frozen at -45°C and lyophilized under 50 μ vacuum with a drying
time of approximately 28 hours. The vials were sealed with split rubber stoppers and capped with
45 plastic screw caps. The freeze dried lysate was reconstituted with 1.2 ml pyrogen-free water for 45
injection and tested with FDA Reference Endotoxin lot EC-2. Sensitivity of enhancing agent treated
lyophilized lysate was 62 pg/ml. A control 40% dilution of LAL without enhancing agent tested before
lyophilization had a sensitivity of 1 ng/ml.

Example VI

- 50 Limulus Lysate Day Pools of a sensitivity equal to or greater than 500 pg/ml *E. coli* endotoxin ICF 50
were combined and prepared as a 40% dilution in 0.025M tris(hydroxymethyl) aminomethane maleate
buffer, pH 7.0, containing in final concentrations 0.02M MgCl₂, 0.01M SrCl₂, 0.01M CaCl₂, and
0.025% ZWITTERGENT™ 3—14. This solution was dispensed in 1.2 ml and 5.2 ml aliquots in 10 ml
vials and frozen at -50°C. The samples were lyophilized under 100 μ vacuum with a drying time of
55 approximately 72 hours. The lyophilized product was reconstituted with 1.2 ml or 5.2 ml pyrogen-free 55
Water for Injection depending on the starting volume of lysate. When tested with *E. coli* endotoxin ICF
following lyophilization, the sensitivity of the enhancing agent treated lyophilized lysate for both
sample sizes was 25 pg/ml in comparison to the control sensitivity of 500 pg/ml.

Claims


1. A process for treating under lysate treating conditions *Limulus* amebocyte lysate having decreased sensitivity to endotoxin due to the presence of an endogenous lysate inhibitor with an enhancing amount of a lysate sensitivity enhancing agent having lysate sensitivity enhancing characteristics to neutralize or partially neutralize said lysate inhibitor thereby increasing the lysate sensitivity to endotoxin, said lysate enhancing agent being selected from the group consisting of
- (I) amphoteric surfactants having the following formulae:



wherein:

- R_1 is an alkylene radical having from 1 to 4 carbon atoms;
 Y and Y' are each (1) hydrogen, (2) lower alkyl or (3) hydroxy lower alkyl;
 R_2 and R_3 are each (1) lower alkyl or (2) hydroxy lower alkyl;
 n is 0 or 1, when n is 0, R_4 is alkyl containing from about 8 to about 18 carbon atoms;
when n is 1, R_4 is an alkylene radical having from 1 to about 6 carbon atoms;
 R_5 is an alkyl containing from about 8 to about 18 carbon atoms;
 M is hydrogen, sodium, potassium or ammonium;


- (II) anionic surfactants having the following formulae:



wherein:

- R_5 , Y and M have the same meaning as set forth above
 R_6 is an alkyl from 8 to 24 carbon atoms
 n_1 is an integer from 1 to 3
 n_2 is 1 or 2
 Ar is phenyl or naphthyl;

(III) cationic surfactants having the following formula:



wherein:

R_5 , Y and Y' have the same meaning as set forth above; and

(IV) nonionic surfactants having the following formula:

...

R R R N-L-O

wherein:

R_5 has the same meaning as set forth above

R_7 and R_8 are each methyl or ethyl; and those nonionic surfactants selected from the group consisting of the condensation product of about 10 to 30 moles of ethylene oxide with the monoester of a hexahydric alcohol containing 6 carbon atoms with the ester group containing 10 to 20 carbon atoms.

2. A process according to Claim 1, wherein the enhancing agent is selected from the group of amphoteric surfactants represented by formula (A) of Claim 1.

3. A process according to Claim 2, wherein the enhancing agent is present in an amount of from about 0.01 to about 0.05% (w/v).

4. A process according to Claim 2 or Claim 3, wherein n is 0, R_4 is tetradecyl, R_2 and R_3 are each methyl, and




is trimethylene.

5. A process according to Claim 2 or Claim 3, wherein n is 1, R_4 is trimethylene, R_2 and R_3 are each methyl and



is



6. A process according to Claim 1, wherein the enhancing agent is selected from the group of amphoteric surfactants represented by formula (D) of Claim 1.

7. A process according to Claim 6, wherein the enhancing agent is present in an amount of from about 0.01 to about 0.05% (w/v).

8. A process according to Claim 6 or Claim 7, wherein n is 1, R_4 is propylene, R_2 and R_3 are each methyl and



is methylene.


9. A process according to Claim 6 or Claim 7, wherein n is 0, R_2 and R_3 are each methyl and




is methylene.

10. A Limulus amebocyte lysate reagent for determining endotoxin comprising a buffered aqueous dispersion of Limulus amebocyte lysate present in an endotoxin determining amount having improved sensitivity to endotoxin and an enhancing amount of a lysate sensitivity enhancing agent having lysate sensitivity enhancing characteristics, said lysate enhancing agent being selected from the group consisting of

(I) amphoteric surfactants having the following formulae:







wherein:

- 5 R_1 is an alkylene radical having from 1 to 4 carbon atoms;
 Y and Y' are each (1) hydrogen, (2) lower alkyl or (3) hydroxy lower alkyl;
 R_2 and R_3 are each (1) lower alkyl or (2) hydroxy lower alkyl;
 n is 0 or 1, when n is 0, R_4 is alkyl containing from about 8 to about 18 carbon atoms;
 when n is 1, R_4 is an alkylene radical having from 1 to about 6 carbon atoms;
 10 R_5 is an alkyl containing from about 8 to about 18 carbon atoms;
 M is hydrogen, sodium, potassium or ammonium;


(II) anionic surfactants having the following formulae:



wherein:

- 15 R_5 , Y and M have the same meaning as set forth above
 R_6 is an alkyl from 8 to 24 carbon atoms
 n_1 is an integer from 1 to 3
 n_2 is 1 or 2
 Ar is phenyl or naphthyl;


20 (III) cationic surfactants having the following formula:



wherein:

R_5 , Y and Y' have the same meaning as set forth above; and

(IV) nonionic surfactants having the following formula:



wherein:

- R_5 has the same meaning as set forth above
 R_7 and R_8 are each methyl or ethyl; and the nonionic surfactants selected from the group
 consisting of the condensation product of about 10 to 30 moles of ethylene oxide with the monoester
 30 of a hexahydric alcohol containing 6 carbon atoms with the ester group containing 10 to 20 carbon
 atoms.

11. A reagent according to Claim 10, wherein additionally are present the cations, Na^+ , Mn^{++} and
 Ca^{++} in lysate sensitivity increasing amounts.

12. A reagent according to Claim 10, wherein additionally are present the cations, Na^+ , Sr^{++} , Ca^{++}
 35 and Mg^{++} in lysate sensitivity increasing amounts.

13. A reagent according to any one of Claims 10 to 12 wherein the Lysate is present in an
 amount to detect from about 0.007 to about 0.050 ng/ml of FDA reference endotoxin EC-2.

14. A reagent according to any one of Claims 10 to 13 wherein the enhancing agent is selected
 from the group of amphoteric surfactants represented by Formula A of Claim 10.

- 40 15. A reagent according to Claim 14, wherein n is 0, R_4 is tetradecyl, R_2 and R_3 are each methyl



is trimethylene.


16. A reagent according to Claim 14, wherein n is 1, R_4 is trimethylene, R_2 and R_3 are each methyl and

5



5

is



17. A reagent according to any one of Claims 10 to 13, wherein the enhancing agent is selected from the group of amphoteric surfactants represented by formula D of Claim 10.

10 18. A reagent according to Claim 17, wherein n is 1, R_4 is propylene, R_2 and R_3 are each methyl and 10



is methylene.

19. A reagent according to Claim 17, wherein n is 0, R_2 and R_3 are each methyl and

15



15

is methylene.

20. A reagent according to any one of Claims 10 to 19 which is lyophilized.

21. A process according to Claim 1 conducted substantially as herein described and exemplified in any one of Examples I to VI.

20 22. A limulus amebocyte lysate reagent for determining endotoxin whenever prepared by a process according to any one of Claims 1 to 9 and 21. 20

23. Limulus amebocyte lysate reagents for determining endotoxin substantially as herein described and exemplified in any one of Examples I to VI.

25 24. In a method for determining endotoxin under endotoxin determining conditions wherein the endotoxin is reacted with a Limulus amebocyte lysate reagent, the improvement comprising utilizing the Limulus amebocyte lysate reagent of any one of Claims 10 to 20, 22 and 23. 25

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.